AGRAMECTIN 1 % + Clor - injectable solution

Composition:
Each ml solution contains : 10 mg Ivermectin and 100 mg Clorsulon .

Activity:
Agramectin Clor 1 % is a combination preparation containing Ivermectine active against parasitic nematodes, arachnids and Clorsulon which is a sulphonamide acting primarily against adult and immature liver flukes.
Ivermectin binds selectively and with high affinity to glutamate-gated chloride ion channels, leading to an increase in the permeability of the cell membrane to chloride ions and finally resulting in hyperpolarization and consequently paralysis and death of the parasite.
Clorsulon is rapidly asborbed into the blood and acts by the inhibition of enzymes in the glycolytic pathway.

Indications:
Treatment and control of gastro-intestinal nematodes, lungworms (Adults and fourth-stage larvae), liver fluke (Fasciola hepatica and F. gigantica; adult stages, eye worms, warbles (parasitic stage), sucking lice and mange mites (scabies) in beef cattle and non-lactating dairy cattle.

Dosage and Administration: For subcutaneous injection.
The general dose is 1 ml per 50 kg bodyweight.

Recommendations and Warnings:
Do not administer to lactating animals and pregant heifers within 60 days of claving.
Avoid water runoff from feedlots to enter lakes and streams.
Dispose of containers by incineration.

Withdrawal time: For meat: 35 days.

Storage: Keep closed in a cool (between 15-25 °C) and dry place, away from sunlight and safe from children.

Packing:
4 x 24 = 96 x 50 ml vials in 1 carton.

FOR VETERINARY USE ONLY