

AGLEVA - water soluble powder



Composition:

Each gram powder contains :
200 mg Levamisole hydrochloride

Indications:

Agleva 20 is indicated in adult and larval stages of gastro-intestinal nematodes and lungworms of cattle, sheep, goats, pigs and poultry.

Agleva 20 is a synthetic broad spectrum anthelmintic and is active against adult and larval stages of gastro-intestinal nematodes and lungworms of cattle, sheep, goats and pigs. After oral or parenteral administration, levamisole first paralyzes the worms and afterwards the worms are expelled in the faeces (within 24 hours after parenteral administration). In therapeutic dose levamisole is non-toxic and no side-effects are observed. Levamisole may be given to pregnant animals or animals in poor condition.

Lungworms (*Dictyocaulus* spp., *Protostrongylus* spp. and *Metastrongylus elongatus* (pigs)). Ascaridae (*Toxocara vitulorum*, *Ascaris suum*).

Trichuridae (*Trichuris ovis*, *Trichuris suis*).

Trichostrongylidae (*Trichostrongylus* spp., *Cooperia* spp., *Ostertagia* spp., *Marshallagia* spp., *Haemonchus* spp., *Nematodirus* spp.).

Strongyloidea (*Oesophagostomum* spp., *Chabertia ovina*, *Bunostomum* spp.). Strongyloidea (*Strongyloides* spp.).

Contra-Indications :

Simultaneous administration with substances having nicotinic like action e.g. pyrantel, morantel.

Dosage and Administrations : For oral administration with the drinking water.

Cattle, sheep, goats and pigs: general dose is 10 mg/kg bodyweight (10 gm powder per 200 kg b.w.).

Poultry: 100 ppm. This is achieved by adding 100 gram powder per 200 L of drinking water

As a rule, one single treatment will be sufficient.

Withdrawal time : 8 days before slaughtering and 3 days for milking.

Storage: Keep closed in a cool (between 15-25 deg C) and dry place, away from sunlight and safe from children..

Packing:

12 x 10 x 100 gm sachet in 1 carton.

12 x 1 kg jar in 1 carton.

FOR VETERINARY USE ONLY