

AGRACOX-O - water soluble coccidiostat powder



Composition:

Each gram powder contains:
Sulfadimerazine sodium: 200 mg
Sulfadiazine sodium: 25 mg
Pyrimethamine : 25 mg
Oxytetracycline HCL: 100 mg
Vitamin A: 15.000 IU
Vitamin K3: 5 mg
Support soluble: q.s. ad 1000 mg

Indications:

Poultry: prevention and treatment of intestinal coccidiosis, diarrhoea, enteritis, Pullorum disease and fowl typhoid in broilers, pullets as well as layers and breeders.

Piglets: prevention and treatment of diarrhoea and enteritis.

Dosage and Administration:

Prevention: 100 grams per 200 litres of drinking water, during 3 consecutive days
Treatment: 100 grams per 100 litres of drinking water, during 3 consecutive days.
Administer during treatment only medicated drinking water.

Recommendations and Warnings:

Medicated drinking water should be used within 24 hours after preparation.

Withdrawal time: 14 days for meat and 4 days for eggs.

Storage: Keep closed in a cool and dry place, away from sunlight and safe from children.

Packing:

12 x 10 sachets of 100 gm sachet in 1 carton.
12 x 1 kg jar in 1 carton.

FOR VETERINARY USE ONLY