



AGRAMINOVIT – oral solution

Composition:

Each ml solution contains:

Vit.A, retinol-palmitate: 10.000 iu	Cysteine: 0.06 mg
Vit.D, cholecalciferol: 3.000 iu	Glutamic acid: 5.94 mg
Vit.E, α -tocopherol: 30 mg	Glycine: 11.26 mg
Vit.B1, thiamine hydrochloride: 3 mg	Histidine: 0.72 mg
Vit.B2, riboflavine sodium ph: 6 mg	Iso-leucine: 0.90 mg
Vit.B6, pyridoxine Hcl: 4 mg	Leucine: 1.98 mg
Vit.B12, cyanocobalamin: 30 μ g	Lysine: 11.62 mg
Vit.C, ascorbic acid: 20 mg	Methionine: 10.36 mg
Vit. K3: 2 mg	Phenylalanine: 1.26 mg
D-Panthenol: 10 mg	Proline: 1.62 mg
Nicotinamide: 40 mg	Serine: 1.44 mg
Biotin: 150 μ g	Threonine: 0.90 mg
Folic acid: 1 mg	Tryptophane: 0.08 mg
Choline chloride: 25 mg	Tyrosine: 0.72 mg
Alanine: 1.44 mg	Valine: 0.90 mg
Arginine: 1.80 mg	Solvents ad: 1 ml
Aspartic acid: 8 mg	

Indications:

Agraminovit is a well balanced combination of essential vitamins and free amino-acids for farm animals and is used for stress conditions, to improve growth and egg-production, improper feed, convalescence and general vitamin and/or amino-acid deficiencies in farm animals.

Dosage and Administration: Orally, via the drinking water.

Poultry and swine: 1 ltr. per 2000 litres of drinking water, during 3-5 consecutive days.

Calves, sheep, goats: 1 ml per 10-20 kg bodyweight, during 3-5 consecutive days.

Cattle, horses: 1 ml per 10-20 kg bodyweight, during 3-5 consecutive days.

Recommendations and Warnings:

Medicated drinking water should be used within 24 hours after preparation.

Storage: Keep closed in a cool and dry place, away from sunlight and safe from children.

Packing:

10 x 1 ltr bottle in 1 carton.

FOR VETERINARY USE ONLY